



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Public health significance of invasive mosquitoes in Europe

Schaffner, Francis ; Medlock, J M ; Van Bortel, W

Abstract: There are currently five invasive *Aedes* mosquito species known to be established in Europe, namely *Aedes albopictus*, *Aedes aegypti*, *Aedes japonicus*, *Aedes atropalpus* and *Aedes koreicus*. *Aedes albopictus* and *Aedes aegypti* are the incriminated vectors in the recent outbreaks of chikungunya and dengue fever in Europe. However, both laboratory experiments and field observations indicate that these invasive mosquitoes have a potential to also transmit other pathogens of public health importance. Increasing travel and pathogen introduction, expansion of vector distribution, and both environmental and climatic changes are likely to raise the risk of pathogen transmission by these invasive *Aedes* mosquitoes. Their vector status and their involvement in pathogen transmission are dynamic processes that shape the future of mosquito-borne disease epidemiology in Europe. Beside vector surveillance, enhanced disease surveillance will enable the early detection of cases and the prompt implementation of control measures.

DOI: <https://doi.org/10.1111/1469-0691.12189>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-90480>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Schaffner, Francis; Medlock, J M; Van Bortel, W (2013). Public health significance of invasive mosquitoes in Europe. *Clinical Microbiology and Infection*, 19(8):685-692.

DOI: <https://doi.org/10.1111/1469-0691.12189>

Public health significance of invasive mosquitoes in Europe

F. Schaffner^{1,2}, J. M. Medlock³ and W. Van Bortel⁴

1) Avia-GIS, Agro-Veterinary Information and Analysis, Zoersel, Belgium, 2) Vector Entomology Unit, Institute of Parasitology, Vetsuisse Faculty, University of Zürich, Zürich, Switzerland, 3) Medical Entomology & Zoonoses Ecology Group, Emergency Response Division, MRA, Health Protection Agency, Porton Down, Salisbury, UK and 4) Emerging & Vector-borne Diseases Programme, European Centre for Disease Prevention and Control, Stockholm, Sweden

Abstract

There are currently five invasive *Aedes* mosquito species known to be established in Europe, namely *Aedes albopictus*, *Aedes aegypti*, *Aedes japonicus*, *Aedes atropalpus* and *Aedes koreicus*. *Aedes albopictus* and *Aedes aegypti* are the incriminated vectors in the recent outbreaks of chikungunya and dengue fever in Europe. However, both laboratory experiments and field observations indicate that these invasive mosquitoes have a potential to also transmit other pathogens of public health importance. Increasing travel and pathogen introduction, expansion of vector distribution, and both environmental and climatic changes are likely to raise the risk of pathogen transmission by these invasive *Aedes* mosquitoes. Their vector status and their involvement in pathogen transmission are dynamic processes that shape the future of mosquito-borne disease epidemiology in Europe. Beside vector surveillance, enhanced disease surveillance will enable the early detection of cases and the prompt implementation of control measures.

Keywords: *Aedes*, chikungunya, dengue, invasive, mosquito, vector species, vector-borne disease

Article published online: 10 April 2013

Clin Microbiol Infect 2013; **19**: 685–692

Corresponding author: F. Schaffner, Avia-GIS, Risschotlei 33, 2980 Zoersel, Belgium
E-mail: fschaffner@avia-gis.be

Invasive Mosquito Species in Europe

Invasive mosquito species are defined by their ability to colonise new territories and to cause or to be likely to cause harm to the economy, environment, or human health. Human actions are the primary means of introduction of invasive species. A considerable increase in the spread of invasive mosquitoes has been observed within Europe since the late 1990s, with the Asian tiger mosquito *Aedes albopictus* (Fig. 1) having continuously expanded its distribution, and several other container-breeding *Aedes* species now being reported from new countries every year (details about successive introductions and spread in Europe are given in [1]). To date, *Ae. albopictus* has colonised almost all Mediterranean countries, whereas the Asian bush mosquito *Aedes japonicus* (Fig. 2) is spreading widely in Central Europe. Two other species, *Aedes atropalpus* and *Aedes koreicus*, have been introduced on several occasions, leading to the establishment of populations at few foci. *Aedes triseriatus* was intercepted at a point of entry,

and its establishment was prevented by the implementation of immediate control measures. Finally, the yellow fever mosquito *Aedes aegypti* (Fig. 3), which had been introduced into Europe during the 17–19th centuries, existed in southern Europe until its disappearance during the 20th century. This species has now returned, having recently established on Madeira as well as around the Black Sea coast (Russia, Abkhazia, Georgia). These invasive mosquito species are well adapted to synanthropic settings where they exploit the abundant sources of host blood, resting places and larval breeding sites (mainly man-made water containers) [2].

Mosquito-Borne Pathogens Transmitted by Invasive Mosquitoes in Europe

Although mosquito-borne diseases (MBDs) represent a lower burden in temperate compared to tropical regions where they pose a substantial impact on the countries' socio-economic



FIG. 1. *Aedes albopictus*, female. Source: F. Schaffner/Avia-GIS.



FIG. 2. *Aedes japonicus*, female. Source: F. Schaffner/Avia-GIS.

development, there have always been both endemic and epidemic autochthonous MBDs in Europe. Concern is now rising as both vectors and pathogens are increasingly being introduced by international travel and trade. Some of these diseases are emerging or are reappearing after a long period of absence. Their occurrence is often associated with changes in ecosystems, human behaviour, and climate [3].

The first notable outbreak of a MBD attributed to invasive mosquitoes in Europe occurred in 2007 in Ravenna, Italy, where more than 200 confirmed chikungunya cases were reported between July and September [4–7]. The index case was most likely a traveller returning from the state of Kerala, India, and the first local case was a relative that he visited [8]. Indeed, the strain that caused the outbreak in Italy was genetically similar to the strains from the Indian subcontinent



FIG. 3. *Aedes aegypti*, female. Source: F. Schaffner/Avia-GIS.

and showed the mutation that is believed to be better adapted to *Ae. albopictus* [6]. The rapid spread of the infection from one index case demonstrated the efficiency of local *Ae. albopictus* populations to transmit chikungunya virus (CHIKV). The second autochthonous transmission of CHIKV in Europe occurred in southern France (Fréjus, department of Var) in 2010. The presumed index case who had returned from Rajasthan, India, sought medical care in late August and lived in the close vicinity of one of the two local cases who developed symptoms in September. The origin of the viral strain circulating in France was confirmed to be from northern India [9].

In the same year, two cases of autochthonous dengue fever occurred in Nice, France. This was the first local transmission of dengue virus (DENV) in Europe, since the major dengue epidemic of 1927–1928 in Greece where *Ae. aegypti* was implicated as the vector [10]. The first patient developed symptoms in late August 2010 and was detected through a routine enhanced surveillance system coordinated by the French authorities in the departments where *Ae. albopictus* had become established. The second case lived near the first case and developed symptoms by mid-September 2010 [11]. Subsequently, Croatia was the second country in Europe with autochthonous transmission of DENV in 2010 [12]. The first case was reported by the German health authorities that notified a laboratory-confirmed case in a German citizen returning from southern Croatia. The patient had spent 2 weeks on the Pelješac peninsula in early August [13]. A second case was reported in late October through active case finding in the same village where the infected tourist had stayed [12]. In addition, further evidence of indigenous

transmission was suggested by positive laboratory results from 14 healthy individuals [12] and by a sero-prevalence survey using a random sample of the population living in the area [14]. Currently, a large outbreak of dengue fever is on-going in the Portuguese Autonomous Region of Madeira [15]. The epidemic started in October 2012 and by early January 2013 more than 2000 cases of dengue fever had been reported, with an additional 70 cases reported among European travellers returning from the island [16]. The presence of *Ae. aegypti* in Madeira has been known since 2005 but this is the first recorded outbreak of dengue on the island.

Since the 2007 outbreak of chikungunya in Italy, it is acknowledged that Europe is vulnerable for transmission of 'tropical' arboviruses, particularly in regions where *Ae. albopictus* or *Ae. aegypti* are present.

Potential Vector Role of Invasive Mosquitoes

Outside Europe, *Ae. albopictus* was implicated in transmission of CHIKV causing large scale disease outbreaks in the Indian Ocean and south Asia since 2005 [17]. Previously, *Ae. albopictus* was responsible for DENV transmission in the 1977–78 Reunion Island epidemic, in an outbreak in Hawaii in 2001–2002 [18] and again in Reunion Island (2004) [19], Gabon (2007) [20], and Mauritius (2009) [21]. *Aedes albopictus* is also a known vector of *Dirofilaria* filarial nematodes, which primarily are parasites of dogs but can also affect humans. Recent evidence of transmission by Italian *Ae. albopictus* populations [22,23] has been linked to an increase in prevalence of human dirofilariosis [24]. Other arboviruses important to human health that have been isolated from field-collected *Ae. albopictus* or for which laboratory transmission has been demonstrated [25] include Eastern equine encephalitis (EEEV) [26,27], Jamestown Canyon virus (JCV) [28], Japanese encephalitis virus (JEV) [25], La Crosse virus (LACV) [29,30], Venezuelan equine encephalitis virus (VEEV) [31,32], and West Nile virus (WNV) [33,34]. Field isolations and experimental infection studies alone, however, simply indicate the potential of a mosquito species to transmit these viruses but cannot prove its epidemiological role. Major dengue fever epidemics due to *Ae. aegypti* occur in the Americas, South East Asia and the western Pacific, and the disease is now endemic in >100 countries worldwide [35]. When this mosquito was widespread in southern Europe, it was responsible for large epidemics of dengue and yellow fever [10,36,37]. *Aedes aegypti* has recently been implicated in outbreaks of chikungunya, notably in Kenya and the Comoros Islands [38]. It is also a highly effective vector of yellow fever virus (YFV) in Africa and South America [39]. Historically, local outbreaks of yellow

fever were reported from several ports in Europe (France, Great Britain, Italy, Portugal, Spain) in the 19th century following importations of the vector and viraemic passengers aboard vessels, with occasional large inland outbreaks where *Ae. aegypti* was established (e.g. Barcelona, Lisbon) [37]. Imported yellow fever cases are occasionally noted across Europe [40–42] with high case-fatality rates. In addition, *Ae. aegypti* was suggested as a vector of Zika virus (ZIKV) owing to virus isolation from field collections [43] and virus transmission under laboratory conditions [44].

Regarding the other invasive species, laboratory competency studies have shown the ability of *Ae. atropalpus* to transmit LACV [45], WNV [46] and other arboviruses causing encephalitis, but its importance as a vector of human pathogens is still not clear [47]. Field-collected *Ae. japonicus* have been found positive for WNV on a number of occasions in the US [48], and laboratory studies showed its high vector competence for WNV [46,49] and also for DENV [50], JEV [51], and LACV [52], whereas it was a moderately efficient vector for CHIKV [50], EEEV [53] and Saint Louis encephalitis virus (SLEV) [54]. However, its role in the transmission of these viruses under natural conditions remains unclear [55]. In North America, *Ae. triseriatus* is a known vector of LACV causing serious disease in humans [56,57], and JCV has repeatedly been isolated from field-collected specimens in the US [58]. It is also suggested to be a possible bridge vector for WNV, based on virus isolations from field collections [59] and vector competence demonstrated under laboratory conditions [60]. Other laboratory studies have shown vector competence for VEEV [61], EEEV, Western equine encephalitis virus (WEEV), DENV, SLEV and YFV [62]. *Aedes koreicus* has been linked to outbreaks of JEV [63] and *Dirofilaria* [64] but is not a confirmed vector of either pathogen.

Invasive Mosquitoes in Europe: A Growing Threat to Public Health?

Pathogen transmission by arthropod vectors depends on multiple factors which define their vectorial capacity C , i.e. the daily rate of future inoculations originating from a currently infective source. In the equation $C = b m a^2 p^t / (-\ln p)$ [65], where b is the vector competence, i.e. the proportion of vectors that develop infective pathogen stages; m the vector density in relation to host density; a the vector's daily blood-feeding rate; p the vector's daily survival rate; t the duration of the pathogen's extrinsic incubation period in days. The vector capacity is specific to a species population in a defined natural context. The vector competence (b) refers to innate factors influencing the ability of a

vector to transmit a pathogen. It is of primary importance, and it can be assessed in the laboratory by experimental infection, indicating the potential for involvement in transmission (Table 1), which is specific to each mosquito species and may vary between populations of a defined species [66,67]. DNA mutation of an arboviral protein can alter the transmission efficiency of a mosquito species, as observed in *Ae. albopictus* for CHIKV where a single amino acid substitution led to a reduction of the extrinsic incubation period

and to an increase in transmission efficiency [68,69]. Besides being competent for a pathogen, a mosquito species needs to both sustain pathogen development under natural temperature ranges (*t*) and have a sufficient daily survival rate (*p*) to transmit the pathogen. Population density (*m*) and vector-host contact rate (*a*) need to be high enough to sustain the transmission cycle and, in cases where the pathogen has non-human reservoir hosts, the mosquitoes must blood-feed on both these animal hosts and humans.

TABLE 1. Main mosquito-borne pathogens that can be transmitted by invasive aedine mosquitoes

Pathogen	Genus	Clinical importance	Disease occurrence	Role of invasive species (confirmed ^a /laboratory ^b)
Chikungunya virus (CHIKV)	Alphavirus	Chikungunya infection can cause fever, myalgia, rash and strong arthralgia which often lasts for months in up to 65% of patients [88]; Clinical manifestations observed during an epidemic on La Reunion included severe hepatitis, severe maternal and foetal disease and meningoencephalitis [89]; There is currently no licensed vaccine against CHIKV [90]	Endemic in Africa and Asia; Frequently imported into Europe	<i>Ae. aegypti</i> <i>Ae. albopictus</i> <i>Ae. japonicus</i>
Eastern equine encephalitis virus (EEEV)	Alphavirus	Severe cases of EEEV infection involving encephalitis begin with the sudden onset of headache, high fever, chills, and vomiting; The illness may then progress into disorientation, seizures, or coma, with fatality rates reaching up to 70% during some epidemics [91]	Americas; No evidence of introduction into Europe to date	<i>Ae. albopictus</i> <i>Ae. triseriatus</i>
Venezuelan equine encephalitis virus (VEEV)	Alphavirus	Largely subclinical, however can cause a nonspecific viral syndrome and rarely neurological complications such as encephalitis, more frequently in children than in adults, including death (fatality rate of 1 to 3%) [91,92]	Endemic in the Americas; No evidence of introduction into Europe to date	<i>Ae. albopictus</i> <i>Ae. triseriatus</i>
Western equine encephalitis virus (WEEV)	Alphavirus	Symptoms range from mild flu-like illness to severe encephalitis, coma and death, with a fatality rate up to 5%, and higher manifestation rates in children and in elderly persons [91]	Endemic in western part of North America; No evidence of introduction into Europe to date	<i>Ae. triseriatus</i>
Jamestown Canyon virus (JCV)	Bunyavirus	Probably most cases are subclinical, and rare human clinical cases show mild febrile illness and at some occasions acute central nervous system infection including meningitis and encephalitis [58,93]	Endemic in North America; No evidence of introduction into Europe to date	<i>Ae. albopictus</i> <i>Ae. triseriatus</i>
Dengue virus (DENV)	Flavivirus	Dengue consists of 4 viral serotypes; The infection causes a flu-like illness which includes fever, intense headache, muscular and joint pain, anorexia, nausea, vomiting and rashes; Sometimes it leads to life-threatening complications including severe bleeding and shock [21,94]; Severe dengue case fatality rates can reach 50% in untreated cases [78]; There is no vaccine available but several candidates are under clinical trials [95]	Endemic in Africa, Americas and Caribbean, Asia; Frequently imported into Europe	<i>Ae. aegypti</i> <i>Ae. albopictus</i> <i>Ae. japonicus</i> <i>Ae. triseriatus</i>
Japanese encephalitis virus (JEV)	Flavivirus	Most cases are subclinical or result in mild symptoms only; However, a small percentage of cases develop encephalitis, with symptoms including sudden onset of headache, high fever, disorientation, coma, tremors and convulsions; It affects primarily children, fatality rate ranges from 10 to 35% and as many as 35% of survivors may have serious life-long neurologic sequelae [92]; Currently the most important mosquito-transmitted arbovirus causing encephalitis in the world, causing 30,000 to 50,000 human cases every year [91]; Vaccines are available	Endemic in Asia; No clear evidence of introduction into Europe to date	<i>Ae. albopictus</i> <i>Ae. japonicus</i> <i>Ae. koreicus</i>
St. Louis encephalitis virus (SLEV)	Flavivirus	Most cases are subclinical; Initial symptoms in clinically apparent cases (less than 1%) include fever, headache, nausea, vomiting, and tiredness; Severe neuroinvasive disease (often involving meningoencephalitis) occurs more commonly in older adults; In rare cases, long-term disability or death can result [96]	Endemic in the Americas; No evidence of introduction into Europe to date	<i>Ae. japonicus</i> <i>Ae. triseriatus</i>
West Nile virus (WNV)	Flavivirus	80% of WNV infection cases are subclinical, and 20% present mild influenza-like illness (fever, headache, body ache); Less than 1% of the cases develop more severe disease, such as encephalitis, meningitis or meningo-encephalitis, mainly in the elderly with fatality rates up to 20% [92]	Worldwide	<i>Ae. albopictus</i> <i>Ae. atropalpus</i> <i>Ae. japonicus</i> <i>Ae. triseriatus</i>
Yellow fever virus (YFV)	Flavivirus	YFV can cause systemic disease including fever, jaundice, haemorrhage and renal failure; Approximately 200,000 cases are reported annually [97]; Symptoms are present in about 1 in 7 of those infected [42] and the mortality rate is 20-50%; There is an effective live-attenuated virus vaccine for YFV which can help to prevent the disease [92]	Endemic in Africa and South America; Occasionally imported into Europe	<i>Ae. aegypti</i> <i>Ae. triseriatus</i>
Zika virus (ZIKV) La Crosse virus (LACV)	Flavivirus Orthobunyavirus	ZIKV can cause headache, rash, malaise and back pain [98] LACV can cause serious neuroinvasive disease in humans (fatality rate of 0.3%); It is the common cause of paediatric arboviral encephalitis in the US with 40 to 170 cases reported annually and 90% of those infected are under the age of 15 [56]; Case numbers, however, are suspected to be underestimated [57]; Prior to WNV spread it was considered one of the most important mosquito-borne diseases in the US [45]; Most human infections occur from July to September [56]	Epidemic in Africa, Asia Endemic in northern America; No evidence of introduction into Europe to date	<i>Ae. aegypti</i> <i>Ae. albopictus</i> <i>Ae. atropalpus</i> <i>Ae. japonicus</i> <i>Ae. triseriatus</i>
<i>Dirofilaria immitis</i> , <i>D. repens</i>	<i>Dirofilaria</i>	In most human infections, larvae are destroyed by the immune system but in some cases, adult worms can develop, resulting in pulmonary or subcutaneous infections [99]	Endemic in southern Europe and the Americas	<i>Ae. albopictus</i> <i>Ae. koreicus</i>

^aConfirmed' refers to evidence of the implication of a species (in bold characters) as vector for a defined pathogen on the field.

^bLaboratory' refers to demonstration of the ability of a species (in regular characters) to transmit a defined pathogen in laboratory experiments (vector competence).

Introduced mosquitoes may affect human health by (i) concurrently harbouring novel pathogens, (ii) transmitting native pathogens, or (iii) transmitting novel pathogens that were independently introduced [70]. The first scenario occurred during the 18th and 19th centuries in Europe with *Ae. aegypti* carrying YFV and DENV, causing disease outbreaks in ports and other places [10,71]. In the second scenario, the transmission rate of an indigenous pathogen may increase because of a high vector capacity of the invasive vector, as observed for *Dirofilaria* transmission in Italy in areas colonised by *Ae. albopictus* [24]. The third scenario appears to be the most common one for disease emergence, and it applies to the recently observed transmission of DENV and CHIKV, with various time periods elapsed between the introduction of the vector and the first outbreak (e.g. first CHIKV transmission occurred 17 years after the establishment of *Ae. albopictus* in Italy but only after 4 years in southern France), depending upon the global context (e.g. frequency and intensity of epidemics in dengue-endemic areas). The first and third scenarios have the potential to lead to large disease outbreaks, in particular because of a limited immunity of individuals in a population.

Considering the increasing frequency of both dengue and chikungunya epidemics worldwide and the movement of viraemic hosts, it is expected that new autochthonous cases of dengue and/or chikungunya fever will occur in the future [14]. It is estimated that 22.5 million travellers arrive in Europe each year, and as many as 185 000 of these could be viraemic for CHIKV [72]. Environmental changes such as land cover change or changes in weather patterns are known to impact vector-borne disease transmission [73]. Climate change could facilitate higher mosquito densities of *Ae. albopictus* or its establishment or spread beyond the current boundaries. It may result in transient summer expansions, or new foci of established populations, at locations where temperature and humidity are more suitable, thus enhancing the transmission potential for CHIKV and DENV in temperate regions [38,74–76]. However, demographic- and anthropogenic-driven environmental changes combined with globalisation and inefficient public health measures are considered the principal driving forces for the emergence and global spread of dengue in the past 40 years [77]. *Aedes albopictus* mosquitoes have the propensity to feed on multiple hosts and, therefore, contribute to the risk of transmission of zoonotic disease agents, whereas the strict anthropophily of *Ae. aegypti* favours transmission of anthroponosis [25]. Indeed, *Ae. albopictus* is considered to play a minor role compared to *Ae. aegypti* in DENV transmission, due to differences in host preferences and to lower vector competence [25,66]. In summary, the presence of *Ae. aegypti*

and *Ae. albopictus* in Europe and the increasing number of overseas travellers may increase the risk of transmission of DENV and CHIKV in Europe [78]. *Aedes japonicus* deserves attention regarding its potential to transmit pathogens, including DENV and CHIKV, owing to high population densities in urban environments, its continuing spread in central Europe, its vector competence for several viruses under laboratory conditions, and its anthropophilic nature [F. Schaffner unpublished data, 50,79–81]. The other invasive species discussed here (*Ae. atropalpus*, *Ae. koreicus*, *Ae. triseriatus*) currently represent a very low risk for public health in Europe, considering their limited distribution. Given the recent developments, surveillance of all invasive aedine mosquitoes is now essential to detect early the presence of these mosquitoes, to assess the risk of MBDs and to prepare for the control of disease outbreaks [2]. The elimination of the species at an early stage of introduction, before it becomes widely established, has been achieved at some foci (e.g. in France [82], Italy [83], and The Netherlands [84,85]). When the species is widely established, the purpose of the control is usually limited to reducing the risk for disease transmission and biting nuisance. A number of control methods are commonly used such as source reduction, pesticide application, biological control, and public education. These methods are often combined as an integrated vector management strategy which so far seems to give the best results [86,87]. However, the control of *Ae. albopictus* has so far proven difficult with the currently available control methods and alternative innovative techniques are needed [1] as well as strategies to prevent the introduction of invasive *Aedes* species. Co-ordination among countries will be essential considering the cross-boundary aspects of the problem. Furthermore, fundamental knowledge on the relevant breeding places, host preferences and longevity of these invasive mosquito species under European conditions as well as on their vector competences for important viruses is required to better target and control these species. Vector status is a dynamic process that in the future could impact significantly on MBD epidemiology. Beside vector surveillance, enhanced disease surveillance will enable the early detection of cases and the prompt implementation of control measures.

Acknowledgements

This review work was conducted within the framework of the European network of medical entomologists and public health experts (VBORNET) under ECDC service contract ECD 2009/018.

Transparency Declaration

The authors declare no conflicts of interest.

References

- Medlock JM, Hansford KM, Schaffner F et al. A review of the invasive mosquitoes in Europe: ecology, public health risks, and control options. *Vector Borne Zoon Dis* 2012; 12: 435–447.
- European Centre for Disease Prevention Control (ECDC). *Guidelines for the surveillance of invasive mosquitoes in Europe*. Technical Report. Stockholm: ECDC, 2012.
- Randolph SE, Rogers DJ. The arrival, establishment and spread of exotic diseases: patterns and predictions. *Nat Rev Microbiol* 2010; 8: 361–371.
- Angelini R, Finarelli AC, Angelini P et al. An outbreak of chikungunya fever in the province of Ravenna, Italy. *Euro Surveill* 2007; 12: pii=3260.
- Angelini R, Finarelli AC, Angelini P et al. Chikungunya in north-eastern Italy: a summing up of the outbreak. *Euro Surveill* 2007; 12: pii=3313.
- Rezza G, Nicoletti L, Angelini R et al. Infection with chikungunya virus in Italy: an outbreak in a temperate region. *Lancet* 2007; 370: 1840–1846.
- Angelini P, Macini P, Finarelli AC et al. Chikungunya epidemic outbreak in Emilia-Romagna (Italy) during summer 2007. *Parassitologia* 2008; 50: 97–98.
- European Centre for Disease Prevention and Control (ECDC). *Chikungunya in Italy – joint ECDC/WHO visit for a European risk assessment. 17–21. Mission Report* Stockholm: ECDC, 2007.
- Grandadam M, Caro V, Plumet S et al. Chikungunya virus, southeastern France. *Emerg Infect Dis* 2011; 17: 910–913.
- Rosen L. Dengue in Greece in 1927 and 1928 and the pathogenesis of dengue hemorrhagic fever: new data and a different conclusion. *Am J Trop Med Hyg* 1986; 35: 642–653.
- La Roche G, Souares Y, Armengaud A et al. First two autochthonous dengue virus infections in metropolitan France, September 2010. *Euro Surveill* 2010; 15: pii=19676.
- Gjenero-Margan I, Aleraj B, Krajcar D et al. Autochthonous dengue fever in Croatia, August–September 2010. *Euro Surveill* 2011; 16: pii=19805.
- Schmidt-Chanasit J, Haditsch M, Schoneberg I, Gunther S, Stark K, Frank C. Dengue virus infection in a traveller returning from Croatia to Germany. *Euro Surveill* 2010; 15: pii=19677.
- European Centre for Disease Prevention and Control (ECDC). *Annual epidemiological report: Reporting on 2009 surveillance data and 2010 epidemic intelligence data*. Surveillance Report. Stockholm: ECDC, 2011.
- Sousa CA, Clairouin M, Seixas G et al. Ongoing outbreak of dengue type 1 in the Autonomous Region of Madeira, Portugal: preliminary report. *Euro Surveill* 2012; 17: pii=20333.
- European Centre for Disease Prevention and Control (ECDC). *Communicable disease threat report: week 2, 6–12 January 2013*. Stockholm: ECDC, 2013. http://ecdc.europa.eu/en/activities/epidemic-intelligence/pages/epidemicintelligence_threatcommunicationcdtr.aspx [Accessed on 26 March 2013].
- Burt FJ, Rolph MS, Rulli NE, Mahalingam S, Heise MT. Chikungunya: a re-emerging virus. *Lancet* 2012; 379: 662–671.
- Effler PV, Pang L, Kitsutani P et al. Dengue fever, Hawaii, 2001–2002. *Emerg Infect Dis* 2005; 11: 742–749.
- Pierre V, Thiria J, Rachou E, Sissoko D, Lassalle C, Renault P. Epidémie de dengue I à la Réunion en 2004. In: *Journées de veille sanitaire 2005*. Paris: Institut de veille sanitaire, 2005; 64.
- Paupy C, Ollomo B, Kamgang B et al. Comparative role of *Aedes albopictus* and *Aedes aegypti* in the emergence of Dengue and Chikungunya in central Africa. *Vector Borne Zoon Dis* 2010; 10: 259–266.
- Ramchurn SK, Moheput K, Goorah SS. An analysis of a short-lived outbreak of dengue fever in Mauritius. *Euro Surveill* 2009; 14: pii=19314.
- Cancrini G, Frangipane di Regalbono A, Ricci I, Tessarin C, Gabrielli S, Pietrobelli M. *Aedes albopictus* is a natural vector of *Dirofilaria immitis* in Italy. *Vet Parasitol* 2003; 118: 195–202.
- Cancrini G, Romi R, Gabrielli S, Toma L, Di Paolo M, Scaramozzino P. First finding of *Dirofilaria repens* in a natural population of *Aedes albopictus*. *Med Vet Entomol* 2003; 17: 448–451.
- Pampiglione S, Rivasi F, Angeli G et al. *Dirofilaria immitis* due to *Dirofilaria repens* in Italy, an emergent zoonosis: report of 60 new cases. *Histopathology* 2001; 38: 344–354.
- Paupy C, Delatte H, Bagny L, Corbel V, Fontenille D. *Aedes albopictus*, an arbovirus vector: from the darkness to the light. *Microbes Infect* 2009; 11: 1177–1185.
- Mitchell CJ. Geographic spread of *Aedes albopictus* and potential for involvement in arbovirus cycles in the Mediterranean basin. *J Vector Ecol* 1995; 20: 44–58.
- Turell MJ, Beaman JR, Neely GW. Experimental transmission of eastern equine encephalitis virus by strains of *Aedes albopictus* and *A. taeniorhynchus* (Diptera: Culicidae). *J Med Entomol* 1994; 31: 287–290.
- Gottfried KL, Gerhardt RR, Nasci RS et al. Temporal abundance, parity, survival rates, and arbovirus isolation of field-collected container-inhabiting mosquitoes in eastern Tennessee. *J Am Mosq Control Assoc* 2002; 18: 164–172.
- Gerhardt RR, Gottfried KL, Apperson CS et al. First isolation of La Crosse virus from naturally infected *Aedes albopictus*. *Emerg Infect Dis* 2001; 7: 807–811.
- Grimstad PR, Kobayashi JF, Zhang MB, Craig GB Jr. Recently introduced *Aedes albopictus* in the United States: potential vector of La Crosse virus (Bunyaviridae: California serogroup). *J Am Mosq Control Assoc* 1989; 5: 422–427.
- Beaman JR, Turell MJ. Transmission of Venezuelan equine encephalomyelitis virus by strains of *Aedes albopictus* (Diptera: Culicidae) collected in North and South America. *J Med Entomol* 1991; 28: 161–164.
- Turell MJ, Beaman JR. Experimental transmission of Venezuelan equine encephalomyelitis virus by a strain of *Aedes albopictus* (Diptera: Culicidae) from New Orleans, Louisiana. *J Med Entomol* 1992; 29: 802–805.
- Holick J, Kyle A, Ferraro W, Delaney RR, Iwaseczko M. Discovery of *Aedes albopictus* infected with West Nile virus in southeastern Pennsylvania. *J Am Mosq Control Assoc* 2002; 18: 131.
- Sardelis MR, Turell MJ, O'Guinn ML, Andre RG, Roberts DR. Vector competence of three North American strains of *Aedes albopictus* for West Nile virus. *J Am Mosq Control Assoc* 2002; 18: 284–299.
- Thomas SJ, Strickman D, Vaughn DW. Dengue epidemiology: virus epidemiology, ecology, and emergence. *Adv Virus Res* 2003; 61: 235–289.
- Eritja R, Escosa R, Lucientes J, Marqués E, Roiz D, Ruiz S. Worldwide invasion of vector mosquitoes: present European distribution and challenges for Spain. *Biol Invasions* 2005; 7: 87–97.
- Aeger JM. Yellow fever in Europe – A general historical review. *Bull US Yellow Fever Inst* 1902; 3: 1–2.
- Gould EA, Higgs S. Impact of climate change and other factors on emerging arbovirus diseases. *Trans R Soc Trop Med Hyg* 2009; 103: 109–121.
- Monath TP. Yellow fever and dengue - the interactions of virus, vector and host in the re-emergence of epidemic disease. *Semin Virol* 1994; 5: 133–145.

40. Colebunders R. Imported case of confirmed yellow fever detected in Belgium. *Euro Surveill* 2001; 5: pii2058.
41. Kiehl W. Suspected case of haemorrhagic fever confirmed as yellow fever in Germany. *Euro Surveill* 1999; 3: pii=1350.
42. Monath TP, Cetron MS. Prevention of yellow fever in persons traveling to the tropics. *Clin Infect Dis* 2002; 34: 1369–1378.
43. Marchette NJ, Garcia R, Rudnick A. Isolation of Zika virus from *Aedes aegypti* mosquitoes in Malaysia. *Am J Trop Med Hyg* 1969; 18: 411–415.
44. Boorman JP, Porterfield JS. A simple technique for infection of mosquitoes with viruses transmission of Zika virus. *Trans R Soc Trop Med Hyg* 1956; 50: 238–242.
45. Freier JE, Beier JC. Oral and transovarial transmission of La Crosse virus by *Aedes atropalpus*. *Am J Trop Med Hyg* 1984; 33: 708–714.
46. Turell MJ, O'Guinn ML, Dohm DJ, Jones JW. Vector competence of North American mosquitoes (Diptera: Culicidae) for West Nile virus. *J Med Entomol* 2001; 38: 130–134.
47. Scholte EJ, Den Hartog W, Braks M, Reusken C, Dik M, Hessels A. First report of a North American invasive mosquito species *Ochlerotatus atropalpus* (Coquillett) in the Netherlands, 2009. *Euro Surveill* 2009; 14: pii=19400.
48. Andreadis TG, Anderson JF, Munstermann LE, Wolfe RJ, Florin DA. Discovery, distribution, and abundance of the newly introduced mosquito *Ochlerotatus japonicus* (Diptera: Culicidae) in Connecticut, USA. *J Med Entomol* 2001; 38: 774–779.
49. Sardelis MR, Turell MJ. *Ochlerotatus j. japonicus* in Frederick County, Maryland: discovery, distribution, and vector competence for West Nile virus. *J Am Mosq Control Assoc* 2001; 17: 137–141.
50. Schaffner F, Vazeille M, Kaufmann C, Failloux A-B, Mathis A. Vector competence of *Aedes japonicus* for chikungunya and dengue viruses. *Eur Mosq Bull* 2011; 29: 141–142.
51. Takashima I, Rosen L. Horizontal and vertical transmission of Japanese encephalitis virus by *Aedes japonicus* (Diptera: Culicidae). *J Med Entomol* 1989; 26: 454–458.
52. Sardelis MR, Turell MJ, Andre RG. Laboratory transmission of La Crosse virus by *Ochlerotatus j. japonicus* (Diptera: Culicidae). *J Med Entomol* 2002; 39: 635–639.
53. Sardelis MR, Dohm DJ, Pagac B, Andre RG, Turell MJ. Experimental transmission of eastern equine encephalitis virus by *Ochlerotatus j. japonicus* (Diptera: Culicidae). *J Med Entomol* 2002; 39: 480–484.
54. Sardelis MR, Turell MJ, Andre RG. Experimental transmission of St. Louis encephalitis virus by *Ochlerotatus j. japonicus*. *J Am Mosq Control Assoc* 2003; 19: 159–162.
55. Versteirt V, Schaffner F, Garros C, Dekoninck W, Coosemans M, Van Bortel W. Introduction and establishment of the exotic mosquito species *Aedes japonicus japonicus* (Diptera: Culicidae) in Belgium. *J Med Entomol* 2009; 46: 1464–1467.
56. Borucki MK, Kempf BJ, Blitvich BJ, Blair CD, Beaty BJ. La Crosse virus: replication in vertebrate and invertebrate hosts. *Microbes Infect* 2002; 4: 341–350.
57. McJunkin JE, de los Reyes EC, Irazuzta JE et al. La Crosse encephalitis in children. *N Engl J Med* 2001; 344: 801–807.
58. Andreadis TG, Anderson JF, Armstrong PM, Main AJ. Isolations of Jamestown Canyon virus (Bunyaviridae: Orthobunyavirus) from field-collected mosquitoes (Diptera: Culicidae) in Connecticut, USA: a ten-year analysis, 1997–2006. *Vector Borne Zoon Dis* 2008; 8: 175–188.
59. Centers for Disease Control and Prevention (CDC). *Mosquito species producing WNV positives by year*. Atlanta, GA: CDC, 2009. <http://www.cdc.gov/ncidod/dvbid/westnile/mosquitospecies.htm> [Accessed on 28 November, 2012].
60. Styer LM, Kent KA, Albright RG, Bennett CJ, Kramer LD, Bernard KA. Mosquitoes inoculate high doses of West Nile virus as they probe and feed on live hosts. *PLoS Pathog* 2007; 3: 1262–1270.
61. Davis MH, Hogge AL Jr, Ferrell JF, Corristan EC. Mosquito transmission of Venezuelan equine encephalomyelitis virus from experimentally infected frogs. *Am J Trop Med Hyg* 1966; 15: 227–230.
62. Freier JE, Grimstad PR. Transmission of dengue virus by orally infected *Aedes triseriatus*. *Am J Trop Med Hyg* 1983; 32: 1429–1434.
63. Miles JA. Some ecological aspects of the problem of arthropod-borne animal viruses in the Western Pacific and South-East Asia regions. *Bull World Health Organ* 1964; 30: 197–210.
64. Feng L-C. The tree hole species of mosquitoes of Peiping, China. *Chin Med J* 1938; S2: 503–525.
65. Scott TW, Takken W. Feeding strategies of anthropophilic mosquitoes result in increased risk of pathogen transmission. *Trends Parasitol* 2012; 28: 114–121.
66. Lambrechts L, Scott TW, Gubler DJ. Consequences of the expanding global distribution of *Aedes albopictus* for dengue virus transmission. *PLoS Negl Trop Dis* 2010; 4: e646.
67. Vazeille M, Rosen L, Mousson L, Failloux A-B. Low oral receptivity for dengue type 2 viruses of *Aedes albopictus* from Southeast Asia compared with that of *Aedes aegypti*. *Am J Trop Med Hyg* 2003; 68: 203–208.
68. Tsatsarkin KA, Vanlandingham DL, McGee CE, Higgs S. A single mutation in chikungunya virus affects vector specificity and epidemic potential. *PLoS Pathog* 2007; 3: e201.
69. Vazeille M, Moutailler S, Coudrier D et al. Two Chikungunya isolates from the outbreak of La Reunion (Indian Ocean) exhibit different patterns of infection in the mosquito, *Aedes albopictus*. *PLoS ONE* 2007; 2: e1168.
70. Juliano SA, Lounibos LP. Ecology of invasive mosquitoes: effects on resident species and on human health. *Ecol Lett* 2005; 8: 558–574.
71. Van Heiningen TW. Contagiousness of yellow fever in the Netherlands between 1820 and 1825. Medical and sanitary aspects of a polymorphic disease. *Hist Sci Med* 2006; 40: 9–22 [In French].
72. Tilston N, Skelly C, Weinstein P. Pan-European Chikungunya surveillance: designing risk stratified surveillance zones. *Int J Health Geogr* 2009; 8: 61.
73. Vanwambeke SO, Lambin EF. Environmental change and vector-borne diseases: The contribution of remote sensing and spatial analyses. In: Verhasselt YLG, ed, *Geographic Medicine, Encyclopedia of Life Support System (EOLSS)*. Oxford: Developed under the Auspices of the UNESCO [http://www.eolss.net], 2006; 202–218.
74. Genchi C, Rinaldi L, Mortarino M, Genchi M, Cringoli G. Climate and *Dirofilaria* infection in Europe. *Vet Parasitol* 2009; 163: 286–292.
75. Weaver SC, Reisen WK. Present and future arboviral threats. *Antiviral Res* 2010; 85: 328–345.
76. Halstead SB. Dengue virus-mosquito interactions. *Annu Rev Entomol* 2008; 53: 273–291.
77. Ooi E-E, Gubler DJ. Global spread of epidemic dengue: the influence of environmental change. *Future Virol* 2009; 4: 571–580.
78. Seyler T, Grandesso F, Le Strat Y, Tarantola A, Depoortere E. Assessing the risk of importing dengue and chikungunya viruses to the European Union. *Epidemics* 2009; 1: 175–184.
79. Kampen H, Zielke D, Werner D. A new focus of *Aedes japonicus japonicus* (Diptera, Culicidae) distribution in western Germany: rapid spread or a further introduction event? *Parasit Vectors* 2012; 5: 284.
80. Schaffner F, Kaufmann C, Hegglin D, Mathis A. The invasive mosquito *Aedes japonicus* in Central Europe. *Med Vet Entomol* 2009; 23: 448–451.
81. Seidel B, Duh D, Nowotny N, Allerberger F. First record of the mosquitoes *Aedes (Ochlerotatus) japonicus japonicus* (Theobald, 1901) in Austria and Slovenia 2011 and for *Aedes (Stegomyia) albopictus* (Skuse, 1895) in Austria 2012. *Entomolog Zeitschrift* 2012; 122: 223–226.
82. Scholte E-J, Schaffner F. Chapter 14. Waiting for the tiger: establishment and spread of the *Aedes albopictus* mosquito in Europe. In: Takken W, Knols BGJ, eds, *Emerging pests and vector-borne diseases in Europe*. Wageningen: Wageningen Academic Publishers; 2007: 241–260.

83. Romi R, Di Luca M, Majori G. Current status of *Aedes albopictus* and *Aedes atropalpus* in Italy. *J Am Mosq Control Assoc* 1999; 15: 425–427.
84. Scholte EJ, Den Hartog WW, Dik M, Schoelitsz B et al. Introduction and control of three invasive mosquito species in the Netherlands. *Euro Surveill* 2010; 15: pii=97110.
85. Scholte EJ, Dik M, Ibañez Justicia A et al. Findings and control of two invasive exotic mosquito species, *Aedes albopictus* and *Ae. atropalpus* (Diptera: Culicidae) in the Netherlands, 2011. *Eur Mosq Bull* 2011; 30: 1–14.
86. World Health Organization (WHO). *Global strategic framework for integrated vector management*. Report No WHO/CDS/CPE/PVC/2004.10. Geneva: WHO, 2004.
87. Abramides GC, Roiz D, Guitart R, Quintana S, Guerrero I, Gimenez N. Effectiveness of a multiple intervention strategy for the control of the tiger mosquito (*Aedes albopictus*) in Spain. *Trans R Soc Trop Med Hyg* 2011; 105: 281–288.
88. Sambri V, Cavrini F, Rossini G, Pierro A, Landini MP. The 2007 epidemic outbreak of Chikungunya virus infection in the Romagna region of Italy: a new perspective for the possible diffusion of tropical diseases in temperate areas? *New Microbiol* 2008; 31: 303–304.
89. Soumahoro MK, Fontenille D, Turbelin C et al. Imported chikungunya virus infection. *Emerg Infect Dis* 2010; 16: 162–163.
90. Moutailler S, Barre H, Vazeille M, Failloux A-B. Recently introduced *Aedes albopictus* in Corsica is competent to chikungunya virus and in a lesser extent to dengue virus. *Trop Med Int Health* 2009; 14: 1105–1109.
91. Pfeffer M, Dobler G. Emergence of zoonotic arboviruses by animal trade and migration. *Parasit Vectors* 2010; 3: 35.
92. Gubler DJ. The global emergence/resurgence of arboviral diseases as public health problems. *Arch Med Res* 2002; 33: 330–342.
93. Centers for Disease Control & Prevention (CDC). Human Jamestown canyon virus infection - Montana, 2009. *Morb Mortal Wkly Rep* 2011; 60: 652–655.
94. World Health Organization (WHO). *Dengue and severe dengue. Fact sheet n°117*. Geneva: WHO, 2012 [updated November 2012]; <http://www.who.int/mediacentre/factsheets/fs117/en/> [Accessed on 24 January, 2013].
95. World Health Organization (WHO). *Global strategy for dengue prevention and control 2012-2020*. Geneva: WHO, 2012.
96. Spinsanti LI, Diaz LA, Glatstein N et al. Human outbreak of St. Louis encephalitis detected in Argentina, 2005. *J Clin Virol* 2008; 42: 27–33.
97. Barnett ED. Yellow fever: epidemiology and prevention. *Clin Infect Dis* 2007; 44: 850–856.
98. Hayes EB. Zika virus outside Africa. *Emerg Infect Dis* 2009; 15: 1347–1350.
99. Kim MK, Kim CH, Yeom BW, Park SH, Choi SY, Choi JS. The first human case of hepatic dirofilariasis. *J Korean Med Sci* 2002; 17: 686–690.